

On the Polynomial of a Path

Robert A. Beezer

Department of Mathematics

University of Illinois

1409 West Green Street

Urbana, Illinois 61801

Submitted by Richard A. Brualdi

ABSTRACT

Let $A(P_n)$ be the adjacency matrix of the path on n vertices. Suppose that $r(\lambda)$ is a polynomial of degree less than n , and consider the matrix $M = r(A(P_n))$. We determine all polynomials for which M is the adjacency matrix of a graph.

1. INTRODUCTION AND PRELIMINARIES

All graphs will be finite, undirected and without loops or multiple edges. For a graph G , the adjacency matrix, $A(G) = (a_{ij})$, is the 0-1 matrix where $a_{ij} = 1$ if and only if v_i and v_j are adjacent in G . The i th distance graph of G , G_i , has the same vertex set as G , and two vertices are adjacent in G_i if and only if they are a distance i apart in G . For several interesting classes of graphs, $A(G_i)$ is a polynomial in $A(G)$. The best known of these are distance-regular graphs [2, p. 132]. In this case, $A(G_i)$ is a polynomial in $A(G)$ of degree i . Moreover, Weichsel [4] has shown that this property characterizes distance-regular graphs.

Thus, it seems appropriate to ask when a polynomial in the adjacency matrix of a graph will yield the adjacency matrix of another graph. In this note we will investigate this question for a class of trees.

If G and H are graphs, and $r(\lambda)$ is a polynomial such that $r(A(G)) = A(H)$, we shall say that H is generated from G by $r(\lambda)$. The adjacency algebra of G , $\mathcal{A}(G)$, is the algebra of all polynomials in $A(G)$. Define the path on n vertices by $VP_n = \{v_i \mid 1 \leq i \leq n\}$ and $EP_n = \{[v_i, v_{i+1}] \mid 1 \leq i \leq n-1\}$. Let $\chi(P_n; \lambda)$ denote the characteristic polynomial of P_n , and define $\chi(P_0; \lambda) = 1$. We will write the entry in position (i, j) of a matrix M as $m_{i,j}$.

2. THE MATRICES D_m

Throughout this section we shall consider that n is given, and we define D_m for $0 \leq m \leq n-1$ by $D_m = \chi(P_m; A(P_n))$.

LEMMA 2.1. For $0 \leq m \leq n-1$, D_m is a 0-1 matrix and $d_{m,i,j} = 1$ if and only if (a) $i-j \equiv m \pmod{2}$, (b) $|i-j| \leq m$, and (c) $m+2 \leq i+j \leq 2n-m$.

Proof. We proceed by induction on m . It is easy to see that the result is true for $m=0$ and $m=1$.

The characteristic polynomial of P_m satisfies the recursion $\chi(P_m; \lambda) = \lambda\chi(P_{m-1}; \lambda) - \chi(P_{m-2}; \lambda)$ [3]. Thus we find

$$D_m = A(P_n)D_{m-1} - D_{m-2}, \quad (2.1)$$

$$d_{m,i,j} = d_{m-1,i-1,j} + d_{m-1,i+1,j} - d_{m-2,i,j}$$

unless $i=1$ or $i=n$; in that case the application of the criteria of Lemma 2.1 will give $d_{m-1,0,j} = d_{m-1,n+1,j} = 0$.

Now we assume that the lemma is true for $m=0, 1, 2, \dots, t-1$ and use (2.1) to determine the entries of D_t . There are several cases to consider.

Case 1: $i-j \not\equiv t \pmod{2}$. In this case the induction hypothesis gives

$$d_{t-1,i-1,j} = d_{t-1,i+1,j} = d_{t-2,i,j} = 0,$$

so (2.1) shows that $d_{t,i,j} = 0$.

Case 2: $|i-j| > t$, $i-j \equiv t \pmod{2}$. When $i-j > t$, $d_{t-1,i-1,j} = d_{t-1,i+1,j} = d_{t-2,i,j} = 0$, so $d_{t,i,j} = 0$. Similarly, when $i-j < -t$, $d_{t,i,j} = 0$.

Case 3: $|i-j| = t$, $t+2 \leq i+j \leq 2n-t$, $i-j \equiv t \pmod{2}$. When $i-j = t$, $d_{t-1,i-1,j} = 1$ and $d_{t-1,i+1,j} = d_{t-2,i,j} = 0$, so $d_{t,i,j} = 1$. When $i-j = -t$, $d_{t,i,j} = 1$.

Case 4: $|i-j| < t$, $t+2 \leq i+j \leq 2n-t$, $i-j \equiv t \pmod{2}$. In this case we find that $d_{t-1,i-1,j} = d_{t-1,i+1,j} = d_{t-2,i,j} = 1$, so $d_{t,i,j} = 1$.

Case 5: $|i-j| < t$, $i+j = t$ or $i+j = 2n-t+2$, $i-j \equiv t \pmod{2}$. When $i+j = t$, $d_{t-1,i-1,j} = 0$ and $d_{t-1,i+1,j} = d_{t-2,i,j} = 1$, so $d_{t,i,j} = 0$. Similarly, when $i+j = 2n-t+2$, $d_{t,i,j} = 0$.

Case 6: $i+j < t$ or $i+j > 2n-t+2$, $i-j \equiv m \pmod{2}$. When $i+j < t$, $d_{t-1,i-1,j} = d_{t-1,i+1,j} = d_{t-2,i,j} = 0$, so $d_{t,i,j} = 0$. When $i+j > 2n-t+2$, $d_{t,i,j} = 0$.

These six cases prove the validity of the lemma for $m = t$, and thus concludes the induction. ■

LEMMA 2.2. *The set $\{D_m | 0 \leq m \leq n-1\}$ is a basis of $\mathcal{A}(P_n)$.*

Proof. Since P_n has diameter $n-1$, the matrices $I_n, A(P_n), A(P_n)^2, \dots, A(P_n)^{n-1}$ are linearly independent [2, p. 12]. Since the dimension of $\mathcal{A}(P_n)$ is at most n , these powers of $A(P_n)$ are a basis of $\mathcal{A}(P_n)$.

Define the linear transformation ψ from $\mathcal{A}(P_n)$ to $\mathcal{A}(P_n)$ by $\psi(A(P_n)^m) = D_m$, $0 \leq m \leq n-1$. Since $\chi(P_m; \lambda)$ is a polynomial of degree m , $D_m \in \mathcal{A}(P_n)$ and ψ has full rank. Thus, the image of the basis $\{A(P_n)^m | 0 \leq m \leq n-1\}$, $\{D_m | 0 \leq m \leq n-1\}$, is a basis of $\mathcal{A}(P_n)$. ■

3. THE MAIN RESULT

Define the integer l by $l = \lfloor n/2 \rfloor$.

THEOREM 3.1. *Suppose that $r(\lambda)$ is a polynomial of degree less than n . Then $r(\lambda)$ generates a nonnull graph from P_n if and only if $r(\lambda) = \chi(P_{2i+1}; \lambda)$ for some i , $0 \leq i \leq l-1$.*

Proof. First, suppose that $r(\lambda) = \chi(P_{2i+1}; \lambda)$ for some i , $0 \leq i \leq l-1$. Then $r(A(P_n)) = D_{2i+1}$. From Lemma 2.1, it is apparent that D_{2i+1} is a symmetric 0-1 matrix with a zero diagonal, and therefore is the adjacency matrix of a graph.

Now, for the converse, assume that $r(\lambda)$ generates a graph from P_n . For convenience, let $M = r(A(P_n))$. Then M is 0-1 matrix, with a zero diagonal. Since $M \in \mathcal{A}(P_n)$, Lemma 2.2 implies that there are numbers a_j , $0 \leq j \leq n-1$, such that $M = \sum_{j=0}^{n-1} a_j D_j$.

Assume that $a_{2i} \neq 0$ for some i , $0 \leq i \leq \lfloor (n-1)/2 \rfloor$. Then let s be the smallest integer such that $a_{2s} \neq 0$. Then,

$$\begin{aligned} m_{s+1, s+1} &= \sum_{j=0}^{n-1} a_j d_{j, s+1, s+1} = \sum_{i=0}^{\lfloor (n-1)/2 \rfloor} a_{2i} d_{2i, s+1, s+1} \\ &= \sum_{i=0}^s a_{2i} d_{2i, s+1, s+1} = a_{2s} d_{2s, s+1, s+1} = a_{2s}. \end{aligned}$$

Since M has a zero diagonal, $0 = m_{s+1, s+1} = a_{2s}$, which contradicts the choice of s . Therefore, $a_{2i} = 0$, $0 \leq i \leq \lfloor (n-1)/2 \rfloor$.

Now, notice that

$$m_{1, 2i+2} = \sum_{k=0}^{l-1} a_{2k+1} d_{2k+1, 1, 2i+2} = a_{2i+1}, \quad 0 \leq i \leq l-1.$$




FIG. 1. Graphs with adjacency matrix D_{2i+1} , $0 \leq i \leq 5$, $n = 12$.

Because M is a 0-1 matrix, $a_{2i+1} \in \{0, 1\}$, $0 \leq i \leq l-1$. Then

$$m_{l+1,l} = \sum_{k=0}^{l-1} a_{2k+1} d_{2k+1,l+1,l} = \sum_{k=0}^{l-1} a_{2k+1}.$$

This relation implies that at most one of a_{2i+1} , $0 \leq i \leq l-1$, is equal to 1. If they are all zero, then $M = 0$, contrary to the hypothesis. Therefore, $M = D_{2i+1}$ for some i , $0 \leq i \leq l-1$.

The minimum polynomial of P_n has degree n , $r(\lambda)$ has degree less than n , and $\chi(P_{2i+1}; A(P_n)) = D_{2i+1} = r(A(P_n))$, so $r(\lambda) = \chi(P_{2i+1}; \lambda)$ for some i , $0 \leq i \leq l-1$. ■

4. CONCLUSION

The graphs which have D_{2i+1} as their adjacency matrix are displayed in Figure 1 for $n = 12$ and $0 \leq i \leq 5$.

A computer study was performed to determine if there are any other situations similar to Theorem 3.1. The only case of interest involved sunsets. A sunset is a path of even length greater than two, with additional vertices adjacent to the central vertex. There exists a nontrivial polynomial which generates a graph from a sunset [1]. It is interesting that the polynomial is a divisor of the characteristic polynomial of a subgraph of the sunset, and the generated graph is isomorphic with the original sunset.

I would like to thank my adviser, Paul M. Weichsel, for his help and encouragement, and the referee for his helpful suggestions.

REFERENCES

1. R. A. Beezer, Polynomials of the adjacency matrix of a graph, Ph.D. Thesis, Univ. of Illinois, Urbana, Ill., 1984.
2. N. L. Biggs, *Algebraic Graph Theory*, Cambridge U.P., Cambridge, 1974.
3. D. M. Cvetkovic, M. Doob, and M. Sachs, *Spectra of Graphs*, Academic, New York, 1980, p. 59.
4. P. M. Weichsel, On distance-regularity in graphs, *J. Combin. Theory Ser. B* 32:156-161 (1982).

Received 20 May 1983; revised 6 March 1984